

# Cardiorespiratory Fitness Predicts Changes in Adiposity in Overweight Hispanic Boys

Courtney E. Byrd-Williams<sup>1</sup>, Gabriel Q. Shaibi<sup>2</sup>, Ping Sun<sup>1</sup>, Christianne J. Lane<sup>1</sup>, Emily E. Ventura<sup>1</sup>, Jaimie N. Davis<sup>1</sup>, Louise A. Kelly<sup>1</sup> and Michael I. Goran<sup>1,3,4</sup>

We have previously shown that cardiorespiratory fitness predicts increasing fat mass during growth in white and African-American youth, but limited data are available examining this issue in Hispanic youth. Study participants were 160 (53% boys) overweight (BMI  $\geq$  85th percentile for age and gender) Hispanic children (mean  $\pm$  s.d. age at baseline =  $11.2 \pm 1.7$  years). Cardiorespiratory fitness, assessed by  $VO_{2max}$ , was measured through a maximal effort treadmill test at baseline. Body composition through dual-energy X-ray absorptiometry and Tanner stage through clinical exam were measured at baseline and annually thereafter for up to 4 years. Linear mixed models were used to examine the gender-specific relationship between  $VO_{2max}$  and increases in adiposity (change in fat mass independent of change in lean tissue mass) over 4 years. The analysis was adjusted for changes in Tanner stage, age, and lean tissue mass. In boys, higher  $VO_{2max}$  at baseline was inversely associated with the rate of increase in adiposity ( $\beta = -0.001$ ,  $P = 0.03$ ); this effect translates to a 15% higher  $VO_{2max}$  at baseline resulting in a 1.38kg lower fat mass gain over 4 years. However,  $VO_{2max}$  was not significantly associated with changes in fat mass in girls ( $\beta = 0.0002$ ,  $P = 0.31$ ). In overweight Hispanic boys, greater cardiorespiratory fitness at baseline was protective against increasing adiposity. In girls however initial cardiorespiratory fitness was not significantly associated with longitudinal changes in adiposity. These results suggest that cardiorespiratory fitness may be an important determinant of changes in adiposity in overweight Hispanic boys but not in girls.

*Obesity* (2008) **16**, 1072–1077. doi:10.1038/oby.2008.16

## INTRODUCTION

The prevalence of overweight in youth is increasing at an alarming rate, and this epidemic is disproportionately affecting ethnic minorities, particularly Hispanic youth (1). The most recent data indicate that ~37% of Mexican-American youth are either at risk for overweight or overweight (2). Of particular public health concern are the negative health consequences associated with adiposity during adolescence, such as risk factors for cardiovascular disease and type 2 diabetes (3). Furthermore, overweight and obesity track into adulthood where negative health consequences extend to include other chronic diseases, including pancreatic and liver cancers (4,5).

There are a myriad of factors that contribute to the trend of increasing adiposity; one of these hypothesized factors is a lack of cardiorespiratory fitness (6,7). Cardiorespiratory fitness is an independent determinant of health across the lifespan (8). In adults, higher cardiorespiratory fitness is protective against cardiovascular disease and all-cause mortality (9) whereas in adolescents, cardiorespiratory fitness has been inversely associated with blood pressure (10), total cholesterol, and

pro-inflammatory markers (11). These adverse associations between fitness and disease risk factors may be a direct function of the relationship between cardiorespiratory fitness and adiposity.

Overweight adolescents exhibit lower cardiorespiratory fitness than their normal weight peers (12). African-American (6) and white youth (6,7) with lower levels of cardiorespiratory fitness experience significantly greater increases in fat mass during growth compared to their more fit counterparts. To our knowledge, the longitudinal association between cardiorespiratory fitness and fat mass has not been examined in Hispanic adolescents.

Arriving at a better understanding of the association between cardiorespiratory fitness and adiposity among Hispanic youth is of particular interest due to the high rates of obesity and increased risk of metabolic disease in this population. Therefore, the purpose of this investigation was to examine whether cardiorespiratory fitness affected the rate of increase in adiposity over time during adolescent growth and whether this was moderated by gender.

<sup>1</sup>Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California, USA; <sup>2</sup>The College of Nursing and Healthcare Innovation, Arizona State University, Phoenix, Arizona, USA; <sup>3</sup>Department of Physiology and Biophysics, Keck School of Medicine, University of Southern California, Los Angeles, California, USA. <sup>4</sup>Present address: Clinical Sciences Centre, University of Southern California, Health Science Campus, Los Angeles, California, USA. Correspondence: Michael I. Goran ([goran@usc.edu](mailto:goran@usc.edu))

Received 4 October 2007; accepted 21 December 2007; published online 28 February 2008. doi:10.1038/oby.2008.16

## METHODS AND PROCEDURES

## Subjects

Study participants in the current investigation consisted of a subgroup of children who are participating in the SOLAR (Study of Latino Adolescents at Risk) Diabetes Project—an ongoing longitudinal study to explore risk factors for the development of type 2 diabetes among Hispanic adolescents. Beginning in 2001, participants between the ages of 8 and 13 were recruited for the study through medical referrals, advertisements, and word of mouth in and around the greater Los Angeles County area. The procedures for the SOLAR Project have been detailed elsewhere (13–15). The current analysis included 160 participants (84 boys and 76 girls) who had completed a baseline measure of  $VO_{2max}$  and had complete data for relevant variables during at least two annual visits. Of the participants excluded from the analysis, 42 were excluded because they failed to reach the criteria for  $VO_{2max}$ , and 39 were excluded because they did not have complete data on all variables used in the current analysis during at least two visits. The participants who were included were not significantly different in percent male, mean age, mean Tanner stage, mean  $VO_{2max}$ , mean total fat mass, or mean total lean tissue mass from the participants who were excluded (all  $P > 0.10$ ). The analysis included a total of 597 observations consisting of 23 students completing two visits, 34 students completing three visits, 66 completing four visits, and 37 students completing five visits. Participants had fewer than five visits for two reasons: participants missed at least one annual visit and the open enrollment design of the study. Participants missing one or more visits were not statistically significantly different than those with complete data in percent male, mean age, Tanner stage,  $VO_{2max}$ , total fat mass, or total lean tissue mass (all  $P > 0.20$ ). Informed consents and assents were obtained from parents and children prior to any testing procedures. The study was approved by the Institutional Review Board of the University of Southern California.

## Methods

**Outpatient screening visit.** On an annual basis, each participant arrived at the University of Southern California General Clinical Research Center at the Los Angeles County Hospital at ~0800 AM following an overnight fast. At the screening visit, a physical examination including the assessment of maturation according to established criteria, Tanner stage (16), was performed by a licensed pediatric health care provider. A detailed medical history was conducted, including parental interview detailing family history of diabetes, ethnicity, and demographics of the participants. Height using wall-mounted stadiometer and weight using a medical balance beam were measured to the nearest 0.1 cm and 0.1 kg, respectively.

Children who met the following screening criteria were invited back for further testing during an inpatient General Clinical Research Center visit: (i) aged 8–13 years at baseline assessment, (ii) BMI  $\geq$  85th percentile for age and gender (at risk for overweight and overweight) defined by the Centers for Disease Control and Prevention standards (17) calculated from the height and weight measurements completed at the General Clinical Research Center outpatient screening, (iii) absence of type 1 and type 2 diabetes defined by the guidelines provided by the American Diabetes Association, (iv) positive family history for type 2 diabetes (sibling, parent, or grandparent), (v) Hispanic ethnicity (all four grandparents of Hispanic descent), (vi) not currently taking medication or diagnosed with a condition known to affect body composition or total energy expenditure or physical activity. For the sake of brevity, the participants will be referred to as overweight instead of at risk for overweight and overweight.

**Inpatient visit.** Participants who met inclusion criteria were admitted to the General Clinical Research Center within 2 weeks after their initial outpatient screening where they completed measurements for cardiorespiratory fitness and body composition. Assessment of cardiorespiratory fitness was collected at baseline assessment only. All other measures were assessed annually for up to 4 years.

**Cardiorespiratory fitness.** To assess cardiorespiratory fitness, participants completed an all-out, progressive, continuous treadmill protocol, which has been described previously (13,14) and is summarized here. The participants walked for 4 min at 0% grade and 4 km/h, after which the treadmill grade was raised to 10%. Each ensuing work level lasted 2 min, during which the grade was increased 2.5%. The speed remained constant until a 22.5% grade was reached at which time the speed was increased by 0.6 km/h until the subject reached exhaustion. Respiratory gases were collected and measured using open circuit spirometry and analyzed on a Med Graphics  $CardiO_2/CP$  Combined Exercise System (MedGraphics, St Paul, MN). Heart rate was measured continuously using a Polar Vantage XL heart rate monitor (Polar, Port Washington, NY). Absolute  $VO_{2max}$  (ml of  $O_2$  per min) was used as a measure of cardiorespiratory fitness. Criteria for  $VO_{2max}$  included participants meeting two of three criteria during a maximal effort treadmill test: (i) respiratory exchange ratio  $>1.0$ , (ii) heart rate  $\geq 195$  beats per min, and (iii) a leveling of  $VO_2$  (defined as an increase of  $O_2$  uptake  $<2$  ml/kg/min with a simultaneously increasing workload).  $VO_{2max}$  was defined as the highest oxygen uptake obtained over the final 1.5 min of the protocol.

**Body composition.** A whole-body dual-energy X-ray absorptiometry scan was performed to assess total body composition, including fat mass, soft lean tissue mass, and percentage body fat, using a Hologic QDR 4500W (Hologic, Bedford, MA).

## Statistical analysis

Descriptive statistics were generated for variables of interest and demographic variables. Students  $t$ -tests were conducted to determine whether there were statistically significant gender differences in baseline characteristics.

For longitudinal analyses, linear mixed-effects models were then used to examine whether baseline  $VO_{2max}$  was associated with total fat mass at age 11 and whether baseline  $VO_{2max}$  was associated with the change in total fat mass as the youth age after adjusting for the changes in total lean tissue mass, Tanner stage, gender, and age. Additionally, an interaction term was included in the mixed-effects regression models to investigate whether gender was an effect modifier of the relationship between cardiorespiratory fitness at baseline and changes in adiposity. Because the interaction term was found to be statistically significant, the data were stratified by gender, and the relationship was re-examined in boys and girls separately. All statistical analyses were performed using Statistical Analysis Software (v9.1; SAS Institute, Cary, NC). The linear mixed-effect analyses were performed using the Mixed Procedure.

Although  $VO_{2max}$  is generally expressed relative to total body mass or fat-free mass (FFM), we chose to express  $VO_{2max}$  in absolute terms (ml/min). There is currently no consensus on the appropriate units of expression when comparing participants with differing body composition, gender, and age. It has previously been shown that the ratio method, which adjusts for body composition by dividing  $VO_{2max}$  values by body mass or FFM, may create spurious results, because it fails to take into account the nonzero intercept (18). It has also been shown that using a regression-based method is a more appropriate approach in normalizing  $VO_{2max}$  data (18), and as a result lean tissue mass was used as a covariate in the model in the current study. Even though the ratio methods were not used in the regression analysis, to facilitate comparison among studies, we reported baseline mean  $VO_{2max}$  relative to FFM (ml/kg FFM/min) and total body mass (ml/kg/min).

## RESULTS

Baseline characteristics for the 160 participants (84 boys and 76 girls) are shown in **Table 1**. At baseline, there were statistically significant gender differences in absolute  $VO_{2max}$  (ml/min),  $VO_{2max}$  divided by body mass (ml/kg/min),  $VO_{2max}$  divided by FFM (ml/kg FFM/min), Tanner stage, and total

**Table 1 Participant characteristics at baseline, mean ± s.d.**

Characteristics	Boys (n = 84)	Girls (n = 76)
Age (years)	11.2 ± 1.6	11.2 ± 1.8
Weight (kg)	62.5 ± 16.8	65.6 ± 20.1
Height (m)	150.7 ± 10.9	149.4 ± 11.2
BMI (kg/m <sup>2</sup> )	27.3 ± 4.6	28.8 ± 5.4
BMI Z-score	2.0 ± 0.4	2.1 ± 0.4
BMI percentile	96.8 ± 3.3	97.2 ± 2.9
Total body fat (kg)	23.1 ± 9.0	26.2 ± 10.3*
Total lean tissue mass (kg)	37.0 ± 9.2	36.9 ± 9.9
Tanner stage (1–5)	1.7 ± 1.0	2.97 ± 1.4***
VO <sub>2max</sub> (ml/min)	2,292.6 ± 574.1	2,043.6 ± 475.0**
VO <sub>2max</sub> (ml/kg/min)	37.5 ± 6.9	32.1 ± 5.1***
VO <sub>2max</sub> (ml/kg FFM/min)	58.4 ± 5.8	52.6 ± 5.5***

FFM, fat-free mass.

For gender: \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001.

**Table 2 Number of participants completing 2, 3, 4, and 5 visits**

	Number of visits				Total
	2	3	4	5	
Boys	14	18	35	17	84
Girls	9	16	31	20	76
Total	23	34	66	37	160


body fat (kg; all *P* < 0.05). The number of boys and girls completing two, three, four, and five annual visits is shown in **Table 2**.

In the linear mixed-effects regression model with the total sample, initial VO<sub>2max</sub> was not associated with changes in total fat mass over age after adjusting for changes in Tanner stage, gender, and lean tissue mass (data not shown;  $\beta = -0.0005$ , *P* = 0.18). **Table 3** shows the effect of the predictor of interest, baseline cardiorespiratory fitness, on total fat mass at age 11 after adjusting for covariates; at baseline when the mean age of the participants was 11 years, each one ml/min change in VO<sub>2max</sub> was marginally associated with a 0.002 kg change in total fat mass after adjusting for covariates (*P* < 0.10). **Table 3** also shows the effect of baseline cardiorespiratory fitness on total fat mass over age after adjusting for covariates; each year, a one ml/min change in VO<sub>2max</sub> was significantly and inversely associated with a 0.001 kg change in total fat mass independent of covariates (*P* < 0.01). When gender was investigated as a potential effect modifier (**Table 3**), it was found to be a significant effect modifier in the relationship between VO<sub>2max</sub> and change in total fat mass after adjusting for changes in Tanner stage and lean tissue mass ( $\beta = 0.002$ , *P* = 0.0007). **Figure 1** is a graphical representation of **Table 3** in which the estimated marginal means of the regression equations were plotted. While holding the covariates constant (intercept = 23.9 kg, Tanner stage = 2, lean tissue mass = 37 kg), the values of gender and VO<sub>2max</sub> were manipulated to illustrate the

**Table 3 Effects of predictors on total fat mass (kg) from linear mixed effects regression (n = 160)**

Predictors	Estimate (s.e.)
Effect of predictors on total fat at age 11	
Intercept	23.95 (0.72) <sup>†</sup>
Gender (male = 0, female = 1)	2.78 (1.11)**
Lean tissue mass (kg)	0.48 (0.5) <sup>†</sup>
Tanner stage (1–5)	-0.46 (0.26)*
Age (years)	-0.28 (0.29)
Baseline cardiorespiratory fitness (VO <sub>2max</sub> )	0.002 (0.001)*
Effect of predictors on total fat mass over age	
Gender (male = 0, female = 1)	1.66 (0.31) <sup>†</sup>
Lean tissue mass (kg)	-0.0002 (0.01)
Tanner stage (1–5)	-0.27 (0.10)***
Baseline cardiorespiratory fitness (VO <sub>2max</sub> )	-0.001 (0.0004)***
Baseline cardiorespiratory fitness (VO <sub>2max</sub> ) × gender	0.002 (0.0005) <sup>†</sup>

\**P* < 0.10, \*\**P* < 0.05, \*\*\**P* < 0.01, <sup>†</sup>*P* < 0.001.



**Figure 1** Estimated marginal means for the effects of cardiorespiratory fitness on fat mass over age in overweight Hispanic boys and girls. Model is adjusted for Tanner stage and lean tissue mass. Value used to represent high fitness was 95th percentile of VO<sub>2max</sub>, and value used to represent low fitness was 5th percentile of VO<sub>2max</sub>. Broken line with boxes represent boys low fitness; solid line with boxes represent boys high fitness; broken line with circles represent girls low fitness; and solid line with circles represent girls high fitness.

gender- and fitness level-specific trajectories of total fat mass as the youth age. The value used to represent high fitness was the 95th percentile of VO<sub>2max</sub>, and the value used to represent low fitness was the 5th percentile of VO<sub>2max</sub>.

Subsequent analyses were stratified by gender, as shown in **Table 4**. Among boys, VO<sub>2max</sub> was a significant predictor of change in total fat mass adjusting for changes in lean tissue mass, Tanner stage, and age ( $\beta = -0.001$ , *P* = 0.03). On the other hand, as also shown in **Table 4**, VO<sub>2max</sub> was not a significant predictor of change in total fat mass in girls after controlling for changes in Tanner stage, lean tissue mass, and age ( $\beta = 0.0005$ , *P* = 0.37).

## DISCUSSION

The primary finding of this investigation was that the longitudinal relationship between initial cardiorespiratory fitness and changes in adiposity in children is different in overweight Hispanic boys and girls. In boys, higher initial cardiorespiratory fitness was associated with less subsequent gain in body

**Table 4** Effects of linear mixed effects regressions on total fat mass (kg) by gender

Predictors	Estimate (s.e.)	
	Boys (n = 84)	Girls (n = 76)
Effect of predictors on total fat at age 11		
Intercept	24.31 (0.78) <sup>†</sup>	25.76 (0.63) <sup>†</sup>
Lean tissue mass (kg)	0.37 (0.06) <sup>†</sup>	0.77 (0.07) <sup>†</sup>
Tanner stage (1–5)	–0.26 (0.43)	–0.61 (0.31) <sup>*</sup>
Age (years)	0.35 (0.36)	0.02 (0.28)
Baseline cardiorespiratory fitness (VO <sub>2max</sub> )	0.001 (0.002)	0.002 (0.002)
Effect of predictors on total fat mass over age		
Lean tissue mass (kg)	0.01 (0.02)	–0.01 (0.02)
Tanner stage	–0.44 (0.17) <sup>***</sup>	–0.01 (0.02)
Baseline cardiorespiratory fitness (VO <sub>2max</sub> )	–0.001 (0.0004) <sup>**</sup>	0.0005 (0.0005)

<sup>\*</sup>P < 0.10, <sup>\*\*</sup>P < 0.05, <sup>\*\*\*</sup>P < 0.01, <sup>†</sup>P < 0.001.

fat. However, initial cardiorespiratory fitness was not significantly associated with subsequent fat mass gain in overweight Hispanic girls. These results suggest that the longitudinal association between cardiorespiratory fitness and obesity in Hispanic youth may be gender specific.

In adolescents, cardiorespiratory fitness, defined as the ability of the cardiovascular and respiratory systems to supply oxygen to skeletal muscles during sustained physical activity, correlates to habitual physical activity, which is defined as the amount a person moves around during the day (12,19,20). Importantly, cardiorespiratory fitness is more strongly associated with moderate-to-vigorous physical activity than total physical activity (12,19). Thus, one potential explanation for our findings is that cardiorespiratory fitness may be acting as a proxy indicator of moderate-to-vigorous physical activity. Whereby, Hispanic boys who engage in greater amounts of moderate-to-vigorous physical activity (as indicated by higher initial cardiorespiratory fitness) may have a more balanced energy equation than nonactive boys resulting in less adiposity in the active boys. In support of this hypothesis, our group previously reported (21) cross-sectionally that Hispanic overweight boys engaged in significantly less moderate-to-vigorous physical activity than nonoverweight boys ( $3.2 \pm 2.1\%$  vs.  $4.8 \pm 2.9\%$  of day spent in moderate-to-vigorous activity, respectively;  $P = 0.01$ ), but we did not find a significant difference in moderate-to-vigorous physical activity between overweight and normal weight Hispanic girls in that study ( $2.9 \pm 1.5\%$  vs.  $2.5 \pm 1.6\%$ ;  $P = 0.33$ ). Although those youth were not followed longitudinally, our previous cross-sectional results coupled with our current longitudinal findings suggest that physical activity/cardiorespiratory fitness and obesity are differentially influenced by gender in Hispanic youth.

There is no clear consensus on whether cardiorespiratory fitness in youth is indicative of habitual moderate-to-vigorous physical activity. According to a review that included >25 studies, which used a myriad of methods to assess

cardiorespiratory capacity, there is only a small-to-moderate relationship between physical activity and cardiorespiratory fitness in adolescents (22). While it is acknowledged that other factors, such as genetics (23), contribute to individual levels of cardiorespiratory fitness, studies examining fitness and physical activity make reasonable arguments for the use of cardiorespiratory fitness as a proxy measure for habitual moderate-to-vigorous physical activity at the group level. Regionally and nationally representative studies in Spain (24) and the United States (12) have reported that adolescents who engaged in more moderate-to-vigorous physical activity had higher values of VO<sub>2max</sub>. Nonetheless, we cannot exclude the possibility that undetermined physiologic or behavioral gender differences in our sample were operational in the observed longitudinal associations between cardiorespiratory fitness and adiposity.

The findings from the current study could help to inform future interventions with overweight Hispanic children. Although cardiorespiratory training intervention studies conducted in healthy, nonobese, normotensive youth tend to have negligible effects on body composition, blood lipids, and blood pressure (25,26), cardiorespiratory training interventions in overweight and obese samples have reported more successes (27,28). Training interventions designed for overweight adolescents may result in a reduction of total body weight, though sometimes inconsistently. Additionally, these intervention studies have reported beneficial changes in insulin sensitivity (27), muscular strength, endothelial function, and cardiorespiratory fitness (27). For example, Rowland *et al.* reported a walking training intervention with sedentary overweight adolescents that resulted in an increase in cardiorespiratory fitness of 9.9% (28). The findings from the current study suggest that if interventions were able to improve the cardiorespiratory fitness of overweight Hispanic boys by as little as 15%, this may result in a lesser accumulation of 1.38 kg of fat mass over 4 years than otherwise would have been observed. Additional studies should be conducted with overweight Hispanic adolescents to examine whether increased cardiorespiratory fitness has beneficial effects on body composition and other chronic disease risk factors.

As observed in other studies (29,30), the baseline values of VO<sub>2max</sub> for girls in the current study were significantly lower than the values for boys. Regardless of how VO<sub>2max</sub> was expressed, the cardiorespiratory fitness of girls was at least 11% lower than boys, and this difference in cardiorespiratory fitness will increase as the boys and girls mature (29). When cardiorespiratory fitness levels of the girls from the current study are compared to a nationally representative sample (12), girls in the current study would fall below the lower 40th percentile for age and gender adjusted fitness levels.

The null finding that initial cardiorespiratory fitness at 11 years is not predictive of changes in adiposity may indicate that energy intake in overweight Hispanic girls rather than energy expenditure may be a more important determinant of increases in adiposity. The poor cardiorespiratory fitness levels may be an indication of insufficient levels of physical activity

necessary to counterbalance higher levels of energy intake to prevent increases in adiposity over time. Even more troubling is that low levels of physical activity are likely to decline even more as the girls progress through adolescence and into adulthood (31) at which point low cardiorespiratory fitness levels and a sedentary lifestyle may result in the development of adiposity-related comorbidities such as type 2 diabetes and cardiovascular disease. Thus, targeting overweight Hispanic girls for physical activity interventions coupled with dietary interventions at a young age may help to improve their body composition and energy balance equations resulting in less adiposity increases over time.

The strengths of this study include a precise measure of body composition using dual-energy X-ray absorptiometry (32), a direct measure of  $VO_{2max}$ , the repeated measures design, and the use of mixed-effects regression modeling. The repeated measures design allowed us to follow the same individuals over time to observe changes in adiposity. The use of linear mixed-effects regression models was a distinct benefit, because it accounts for the violation of the assumption of independence across repeated measures and is able to include cases with missing waves of data allowing for a larger sample size to be included in the analysis.

Notwithstanding these strengths, we acknowledge several limitations of the study that may impede the generalizability of the findings. The current study consisted only of overweight Hispanic adolescents with a family history of diabetes. The possibility of gender moderating the relationship between initial cardiorespiratory fitness and changes in adiposity in youths of other ethnicities has not been addressed in this investigation, and as a result, no conclusions should be drawn about the moderating effects of gender in other ethnicities. Also, owing to the homogeneity of the sample, it may not be appropriate to generalize these findings to Hispanic adolescents with a different overweight status or a different family history of disease. Despite the potential limitations of the homogenous sample, we believe the investigation of increasing adiposity in these already overweight adolescents has scientific merit, because the overweight status and ethnicity of these youth increase their risk for chronic diseases, such as type 2 diabetes and cardiovascular disease.

Another limitation is that the current study did not provide repeated measures of dietary data or cardiorespiratory fitness. Dietary data would have allowed for the influence of energy intake on adiposity to be examined. Additionally, because cardiorespiratory fitness was only assessed at baseline, the change in  $VO_{2max}$  and its effect on change in fat mass could not be examined. Future longitudinal studies that include repeated measures of  $VO_{2max}$ , dietary intake and adiposity are warranted. Investigating the association between the change in  $VO_{2max}$  and changes in fat mass independent of energy intake would provide further information on how cardiorespiratory fitness may influence adiposity.

In summary, these findings suggest the relationship between cardiorespiratory fitness and subsequent adiposity may differ by gender suggesting that gender-specific interventions

targeting changes in adiposity may be more effective. To further explain this relationship, additional intervention studies that manipulate cardiorespiratory fitness to examine the effects on adiposity in overweight Hispanic youth are needed, as are longitudinal studies that include direct measures of cardiorespiratory fitness, objective measures of physical activity, and measures of dietary intake.

#### ACKNOWLEDGMENTS

We thank study coordinators, Christina Ayala, and the nursing staff at the University of Southern California General Clinical Research Center. Additionally, we are grateful to our study participants and their families for their involvement. This study was supported by the National Institutes of Health (R01 DK 59211) and National Cancer Institute (Cancer Control and Epidemiology Research Training Grant, T32 CA 09492 and the University of Southern California Center for Transdisciplinary Research on Energetics and Cancer, U54 CA 116848).

#### DISCLOSURE

The authors declared no conflict of interest.

© 2008 The Obesity Society

#### REFERENCES

1. Strauss RS, Pollack HA. Epidemic increase in childhood overweight, 1986–1998. *JAMA* 2001;286:2845–2848.
2. Ogden CL, Carroll MD, Curtin LR *et al*. Prevalence of overweight and obesity in the United States, 1999–2004. *JAMA* 2006;295:1549–1555.
3. Ogden CL, Yanovski SZ, Carroll MD, Flegal KM. The epidemiology of obesity. *Gastroenterology* 2007;132:2087–2102.
4. McMillan DC, Sattar N, Lean M, McArdle CS. Obesity and cancer. *BMJ* 2006;333:1109–1111.
5. Serdula MK, Ivery D, Coates RJ *et al*. Do obese children become obese adults? A review of the literature. *Prev Med* 1993;22:167–177.
6. Johnson MS, Figueroa-Colon R, Herd SL *et al*. Aerobic fitness, not energy expenditure, influences subsequent increase in adiposity in black and white children. *Pediatrics* 2000;106:E50.
7. Koutedakis Y, Bouziotas C, Flouris AD, Nelson PN. Longitudinal modeling of adiposity in periadolescent Greek schoolchildren. *Med Sci Sports Exerc* 2005;37:2070–2074.
8. Ferreira I, Twisk JW, van Mechelen W, Kemper HC, Stehouwer CD. Development of fatness, fitness, and lifestyle from adolescence to the age of 36 years: determinants of the metabolic syndrome in young adults: the amsterdam growth and health longitudinal study. *Arch Intern Med* 2005;165:42–48.
9. Wei M, Kampert JB, Barlow CE *et al*. Relationship between low cardiorespiratory fitness and mortality in normal-weight, overweight, and obese men. *JAMA* 1999;282:1547–1553.
10. Carnethon MR, Gulati M, Greenland P. Prevalence and cardiovascular disease correlates of low cardiorespiratory fitness in adolescents and adults. *JAMA* 2005;294:2981–2988.
11. Barbeau P, Litaker MS, Woods KF *et al*. Hemostatic and inflammatory markers in obese youths: effects of exercise and adiposity. *J Pediatr* 2002;141:415–420.
12. Pate RR, Wang CY, Dowda M, Farrell SW, O'Neill JR. Cardiorespiratory fitness levels among US youth 12 to 19 years of age: findings from the 1999–2002 National Health and Nutrition Examination Survey. *Arch Pediatr Adolesc Med* 2006;160:1005–1012.
13. Ball GD, Shaibi GQ, Cruz ML *et al*. Insulin sensitivity, cardiorespiratory fitness, and physical activity in overweight Hispanic youth. *Obes Res* 2004;12:77–85.
14. Shaibi GQ, Cruz ML, Ball GD *et al*. Cardiovascular fitness and the metabolic syndrome in overweight latino youths. *Med Sci Sports Exerc* 2005;37:922–928.
15. Goran MI, Bergman RN, Avilla Q *et al*. Impaired glucose tolerance and reduced beta-cell function in overweight Latino children with a positive family history of type 2 diabetes. *J Clin Endocrinol Metab* 2004;89:207–212.
16. Tanner JM. Growth and maturation during adolescence. *Nutr Rev* 1981;39:43–55.

17. Ogden CL, Flegal KM, Carroll MD, Johnson CL. Prevalence and trends in overweight among US children and adolescents, 1999–2000. *JAMA* 2002;288:1728–1732.
18. Toth MJ, Goran MI, Ades PA, Howard DB, Poehlman ET. Examination of data normalization procedures for expressing peak VO<sub>2</sub> data. *J Appl Physiol* 1993;75:2288–2292.
19. Dencker M, Thorsson O, Karlsson MK *et al*. Daily physical activity and its relation to aerobic fitness in children aged 8–11 years. *Eur J Appl Physiol* 2006;96:587–592.
20. Rowlands AV, Eston RG, Ingledew DK. Relationship between activity levels, aerobic fitness, and body fat in 8- to 10-yr-old children. *J Appl Physiol* 1999;86:1428–1435.
21. Byrd-Williams C, Kelly LA, Davis JN, Spruijt-Metz D, Goran MI. Influence of gender, BMI and Hispanic ethnicity on physical activity in children. *Int J Pediatr Obes* 2007;2:159–166.
22. Morrow JR, Freedson PS. Relationship between habitual physical activity and aerobic fitness in adolescents. *Pediatr Exerc Sci* 1994;6:315–329.
23. Bouchard C, Rankinen T. Individual differences in response to regular physical activity. *Med Sci Sports Exerc* 2001;33(Suppl 1):S446–S451;discussion S452–S453.
24. Ara I, Moreno LA, Leiva MT, Gutin B, Casajus JA. Adiposity, physical activity, and physical fitness among children from Aragon, Spain. *Obesity (Silver Spring)* 2007;15:1918–1924.
25. Rowland TW. The role of physical activity and fitness in children in the prevention of adult cardiovascular disease. *Prog Pediatr Cardiol* 2001;12:199–203.
26. Rowland TW, Martel L, Vanderburgh P, Manos T, Charkoudian N. The influence of short-term aerobic training on blood lipids in healthy 10–12 year old children. *Int J Sports Med* 1996;17:487–492.
27. Nassis GP, Papatakou K, Skenderi K *et al*. Aerobic exercise training improves insulin sensitivity without changes in body weight, body fat, adiponectin, and inflammatory markers in overweight and obese girls. *Metabolism* 2005;54:1472–1479.
28. Rowland TW, Varzeas MR, Walsh CA. Aerobic responses to walking training in sedentary adolescents. *J Adolesc Health* 1991;12:30–34.
29. Armstrong N, Welsman JR, Nevill AM, Kirby BJ. Modeling growth and maturation changes in peak oxygen uptake in 11–13 yr olds. *J Appl Physiol* 1999;87:2230–2236.
30. Armstrong N, Welsman JR. Peak oxygen uptake in relation to growth and maturation in 11- to 17-year-old humans. *Eur J Appl Physiol* 2001;85:546–551.
31. Kimm SY, Glynn NW, Kriska AM *et al*. Decline in physical activity in black girls and white girls during adolescence. *N Engl J Med* 2002;347:709–715.
32. Goran MI. Measurement issues related to studies of childhood obesity: assessment of body composition, body fat distribution, physical activity, and food intake. *Pediatrics* 1998;101:505–518.